



E-mail :info@kistmcth.edu.np | www.kistmcth.edu.np

Journal of KIST Medical College

Accuracy of Prediction of Birth Weight by Fetal Ultrasound

Sumnima Acharya¹, Awadhesh Tiwari¹

¹Radiology Department, Lumbini Medical College, Palpa.

ABSTRACT

Introduction: Fetal weight estimation using ultrasonography (USG) is beneficial for the better fetomaternal outcome. This study was done to determine the accuracy of prediction of birth weight by fetal ultrasound.

Methods: This was a cross-sectional study carried out in the Department of Radio diagnosis, Lumbini Medical College and Teaching Hospital (LMCTH), Palpa, Nepal from 1st June to 31st December 2018. Fetal weight was calculated by USG in 325 women using Hadlock's formula and correlated with birth weight.

Results: Our study showed that fetal ultrasound using Hadlock's formula has error in estimation of fetal weight by 189gm (SD: 111 gm). In 91.3% of the cases, there was an error of estimation by less than 10% compared to actual weight.

Conclusion: It can be concluded that sonographic estimation of birth weight can be recommended to yield a better prediction of birth weight and to further evaluate fetal well-being.

Keywords: Fetal weight; Fetal ultrasound; Hadlock's formula.

Citation: Acharya S, Tiwari A. Accuracy of prediction of birth weight by fetal ultrasound. JKISTMC 2020;2(2)4: 11-14.

Correspondence

Dr. Sumnima Acharya

Assistant Professor, Radiology Department

Lumbini Medical College, Palpa.

Email: doctorsumnima@gmail.com,

ORCID ID: 0000-0001-8612-671X

Conflict of interest: None

Source of support: None

Article info

Received: 13 March, 2020.

Accepted: 29 May, 2020.

Published: 31 July, 2020.

Copyright

JKISTMC applies the Creative Commons Attribution- Non Commercial 4.0 International License (CC BY) to all works we publish. Under the CC BY license, authors retain ownership of the copyright for their article, but authors allow anyone to download, reuse, reprint, distribute, and/or copy articles in JKISTMC, so long as the original authors and source are cited.



INTRODUCTION

Fetal weight estimation using ultrasonography is useful in timing and mode of delivery of a pregnancy, which is beneficial for the better fetomaternal outcome.¹ There are basically three groups of birth weights that are: the low birth weight, the normal birth weight, and the macrosomic babies.² An Increased risk of newborn complications are associated with extremes of birth weight.³ So recognition of these abnormalities i.e fetal growth restriction (FGR) and large for gestational age (LGA) fetuses are important.⁴ Intrauterine growth restriction (IUGR) or fetal growth restriction (FGR) is defined as an estimated fetal weight (EFW)/abdominal circumference (AC) at one point in time during pregnancy being below 3rd percentile or EFW/AC below the 10th percentile for gestational age with deranged Doppler parameters.⁵ Estimation of birth weight is very important for planning of delivery of very low birth weight baby, the route of delivery and for prenatal counselling.⁶ Prematurity and low birth weights determine the neonatal survival and is an important factor for determining the perinatal morbidity and mortality.⁷ LGA fetuses are those with a birth weight greater than the 90th percentile or >4000gm and are at risk of limb injuries during normal delivery leading to increased cesarean rate.^{4,8}

Estimation of fetal weight has evolved from physical examination to fetal ultrasound. Hence accuracy of fetal weight estimation has increased significantly.^{9,10} There are multiple formulae for the birth weight estimation using ultrasound.¹¹⁻¹⁷ Hadlock's formula is commonly used in Nepal for the estimation of fetal weight using ultrasonography.¹⁸ Our study is directed to see the accuracy of fetal ultrasound in estimating fetal weight in the pregnant patients visiting our hospital.

METHODS

It was an observational, cross-sectional study carried out in the Department of Radio diagnosis, Lumbini Medical College and Teaching Hospital (LMCTH), Palpa, Nepal. The study was conducted after ethical approval from the Institutional Review Committee of our hospital. The study was conducted over a period of six months from 1st June to 31st December 2018. Hospital records (Patient's file sent to radiology department for USG) were reviewed from patients who were sent from Obstetrics and gynecology OPD/

ward with full term pregnancy who came for delivery. Exclusion criteria included multiple pregnancy, preterm birth, intrauterine growth retardation and delivery done more than seven days after USG. Ultrasound examinations were performed by experienced radiologists using standard techniques. Hadlock's formula was used to calculate fetal weight. Observations during the study of each subjects were recorded in an individual case proforma. The case proforma contained all informations regarding the admission details, general particulars like: name, age, Last menstrual period(LMP), Average gestational age (AGA) by date of ultrasound, USG findings including Femoral length(FL),Biparietal diameter(BPD),Head circumference(HC),Abdominal circumference(AC),AGA by USG, Estimated fetal weight(EFW),Amniotic fluid index (AFI),Birth weight of new born baby, date of delivery, mode of delivery. Some informations were recorded in department of radio diagnosis at the time of USG and remaining relevant informations like date, time, mode of delivery, membrane rupture time, and birth weight of new born baby were recorded from discharge summary. Eventually, USG estimated fetal weight were compared with the birth weight of new born baby. We collected the data in Microsoft Excel 2007 and imported it to Statistical Package for the Social Science (SPSSTM), version 16, for the statistical analysis .The absolute error in birth weight prediction was calculated as the absolute value of the difference between the predicted and actual birth weight. The negative values indicate an underestimation of birth weight and positive values indicated overestimation of birth weight. The absolute percent error was calculated by dividing the absolute error in birth weight prediction by the actual birth weight multiplied by 100. Mean error was calculated. Level of significance was set at $p < 0.05$.

RESULTS

The study included 325 patients. The gestational age was between 37 weeks and 42.4 weeks. The age range of patients was between 15-41 years, with a mean of 24.6 years. The range of actual birth weight was between 1.93-4.51 kg with a mean of 2.97 ± 0.424 kg. The mean error in the estimation of birth weight was 189gm (SD: 111 gm).In average, ultrasound overestimate the birth weight by 157 gm (SD: 154gm) and underestimate by 154gm (SD: 108gm). In 49.53% of the cases, fetal ultrasound underestimated the birth

weight and in 50.46% overestimated the birth weight. Twenty eight (8.6%) out of 325 estimates were more than 10% from the actual weight and 91.3% estimates less than 10% from actual birth weight.

Table1. Maternal and Infant Demographics

Characteristics	Mean(Range)
Maternal age(in yrs)	24.6(15-41)
Gestational age at delivery(wks)	39wks5days(37-42.4)
Actual Birth weight(kg)±SD	2.99±0.424(1.93-4.51)
Estimated birth weight(kg)±SD	2.97±0.407(2.08-4.28)

Table 2. Mean Error in birth weight prediction

Characteristics	Mean(kg)±SD
Overestimate	0.157±0.154
Underestimate	0.154±0.108
Absolute	0.189±0.111
Estimate error>10% of ABW-28(8.6%) and <10% of ABW-297(91.3%).	

Table 3. Error estimation

Characteristic	Number(percentage)
Overestimate	164(50.46%)
Underestimate	161(49.53%)

DISCUSSION

For reducing perinatal morbidity and mortality, accurate estimation of antenatal fetal weight is necessary.¹⁹ Many studies have been undertaken to find out the accurate methods of estimation of fetal weight .It includes clinical and ultrasound estimations. Clinical method involves estimation of uterine height and abdominal girth measured at the level of umbilicus.¹⁸ Fetal weight can be measured via ultrasound using abdominal circumference (AC) alone ¹²,AC and biparietal diameter(BPD) ¹³ ,AC ,BPD and femur length.¹⁵ Up to 10% of actual birth weight measured antenatally using ultrasound is considered acceptable.¹⁸

Antenatal birth weight prediction by Parvathavarthini et al was accurate in around 67% of cases. The mean weight of the 100 babies monitored in the study was 2984 grams, ranging from 2000 to 4500gm. Percentage error was least with USG which is almost comparable to our findings.¹⁹

Bajracharya J et al conducted a study on accuracy of prediction of birth weight by fetal USG and found that gestational age was between 37 weeks and 42 weeks. The age range of patients was between 18-40 years, with a mean of 25.51 years. The range of actual birth weight was between 2.11-4.9 kg with a mean of 3.07. The mean error in the estimation of birth weight was 290gm (CI: 250-330 gm).In 56% of the cases, fetal ultrasound overestimated the birth weight and in 36.67% it underestimated the birth weight. In average, ultrasound overestimated by 370 gm (CI: 320-420 gm) and underestimated the birth weight by 220 gm (CI: 160-280gm).¹ Most of the above findings are relatable to our study.

Most of the studies showed that about 75% of the estimates are within 10% of the actual weight.^{6,10} and other studies also showed high correlation between EFW and birth weight^{2,11} and this study shows 91.3% estimates less than 10% from actual birth weight. So the ultrasound method is generally a better predictor of the actual birth weight than the clinical method, and thus should be used in estimating the actual birth weight when accessible.

We also need to keep in mind that ultrasound measurements are operator dependent. There can be high percentage of error during estimation of fetal weight as it is operator dependent. ¹⁸ There is a learning curve for ultrasonographic estimates of fetal weight, with a significant decrease in the percent error seen with advancing training and experience.⁹

CONCLUSION

From the results of the present study it can be concluded that sonographic estimation of birth weight can be recommended to yield a better prediction of birth weight and to further evaluate fetal well-being.

REFERENCES

1. Milner J, Arezina J. The accuracy of ultrasound estimation of fetal weight in comparison to birth weight: A systematic review. Ultrasound [Internet]. 2018 Feb 1 [cited 2019 Oct 21];26(1):32–41.

2. Ugwu EO, Udealor PC, Dim CC, Obi SN, Ozumba BC, Okeke DO, et al. Accuracy of clinical and ultrasound estimation of fetal weight in predicting actual birth weight in Enugu, Southeastern Nigeria. *Niger J Clin Pract.* 2014 Jun;17(3):270–5.
3. Jolly MC, Sebire NJ, Harris JP, Regan L, Robinson S. Risk factors for macrosomia and its clinical consequences: a study of 350,311 pregnancies. *Eur J Obstet Gynecol Reprod Biol.* 2003 Nov 10;111(1):9–14.
4. Haragan AF, Hulse TC, Hawk AF, Newman RB, Chang EY. Diagnostic accuracy of fundal height and handheld ultrasound-measured abdominal circumference to screen for fetal growth abnormalities. *Am J Obstet Gynecol* [Internet]. 2015 Jun [cited 2019 Oct 27];212(6):820.e1-820.e8.
5. Weerakkody Y. Intrauterine growth restriction. *Radiology Reference Article* [Radiopaedia.2019 Oct 27
6. Shittu AS, Kuti O, Orji EO, Makinde NO, Ogunniyi SO, Ayoola OO, et al. Clinical versus sonographic estimation of foetal weight in southwest Nigeria. *J Health Popul Nutr.* 2007 Mar;25(1):14–23.
7. Adimora GN, Odetunde IO. Perinatal mortality in University of Nigeria Teaching Hospital (UNTH) Enugu at the end of the last millennium. *Niger J Clin Pract.* 2007 Mar;10(1):19–23.
8. Ezegwui HU, Ikeako LC, Egbuji C. Fetal macrosomia: Obstetric outcome of 311 cases in UNTH, Enugu, Nigeria. *Niger J Clin Pract* 2019 ;14(3):322.
9. Predanic M, Cho A, Ingrid F, Pellettieri J. Ultrasonographic Estimation of Fetal Weight. *J Ultrasound Med* 2002 ;21(5):495–500.
10. Harlev A, Walfisch A, Bar-David J, HersHKovitz R, Friger M, Hallak M. Maternal estimation of fetal weight as a complementary method of fetal weight assessment: a prospective clinical trial. *J Reprod Med.* 2006 Jul;51(7):515–20.
11. Ben-Haroush A, Yogev Y, Bar J, Mashlach R, Kaplan B, Hod M, et al. Accuracy of sonographically estimated fetal weight in 840 women with different pregnancy complications prior to induction of labor. *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol.* 2004 Feb;23(2):172–6.
12. Campbell S, Wilkin D. Ultrasonic measurement of fetal abdomen circumference in the estimation of fetal weight. *Br J Obstet Gynaecol.* 1975 Sep;82(9):689–97.
13. Campbell S, Thoms A. Ultrasound measurement of the fetal head to abdomen circumference ratio in the assessment of growth retardation. *Br J Obstet Gynaecol.* 1977 Mar;84(3):165–74.
14. Chauhan SP, Hendrix NW, Magann EF, Morrison JC, Kenney SP, Devoe LD. Limitations of clinical and sonographic estimates of birth weight: experience with 1034 parturients. *Obstet Gynecol.* 1998 Jan;91(1):72–7.
15. Hadlock FP, Harrist RB, Carpenter RJ, Deter RL, Park SK. Sonographic estimation of fetal weight. The value of femur length in addition to head and abdomen measurements. *Radiology.* 1984 Feb;150(2):535–40.
16. Jordaan HVF. Estimation of fetal weight by ultrasound. *J Clin Ultrasound* 1983;11(2):59–66.
17. Nzeh DA, Rimmer S, Moore WM, Hunt L. Prediction of birthweight by fetal ultrasound biometry. *Br J Radiol.* 1992 Nov;65(779):987–9.
18. Bajracharya J, Shrestha NS, Karki C. Accuracy of prediction of birth weight by fetal ultrasound. *Kathmandu Univ Med J KUMJ.* 2012 Jun;10(38):74–6.
19. K P, C S, Prasad G. Comparative study of various methods of fetal weight estimation at term pregnancy in a tertiary hospital in Kanchipuram, Tamil Nadu, India. *Int J Reprod Contracept Obstet Gynecol* 2018 Mar 27 ;7(4):1602–7.